



Sustainable Application of NATIVO 300 SC Fungicide (Tebuconazole 200 g/L +Trifloxystrobin 100 g/L) in the Control of Brown Spot (*Bipolaris oryzae*) in Strict Rainfed Rice Cultivation in Côte d'Ivoire

OUATTARA Adama ^{a*}, GUEU Kouato Richmond ^b,
BOUET Alphonse ^a and SORO Kafana ^c

^a Centre National de Recherche Agronomique (CNRA), Abidjan, Km 17, Route de Dabou, 01 B.P. 1470 Abidjan 01, Station de Recherche BP 440 Man, Côte d'Ivoire.

^b Université Nangui Abrogoua 08 BP 109 Abidjan 08, Côte d'Ivoire.

^c Centre de Recherche en Ecologie, Université Nangui Abrogoua 08 BP 109 Abidjan 08, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i113099>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125337>

Original Research Article

Received: 19/09/2024

Accepted: 22/10/2024

Published: 02/12/2024

*Corresponding author: E-mail: adamaouattara85@gmail.com, adama.ouattara@cnra.ci;

ABSTRACT

Rice brown spot is a devastating disease caused by *Bipolaris oryzae*. Various fungicide formulations are used against this phytomycosis, which occurs in all ecologies in Côte d'Ivoire. This study aimed to rationalize the application of a fungicide effective against brown spot in strict rainfed rice cultivation. The experimental design was a Fisher block with three replications. It was carried out in strict rainfed ecology at Biankouma between July and November 2023. The fungicide used was NATIVO 300 SC (Tebuconazole 200 g/L + Trifloxstrobil 100 g/L). Five different treatments were tested (T0: no fungicide treatment; T1: NATIVO 300 SC at a dose of 350 ml.ha⁻¹, every 10 days interval from the initial appearance of brown spot symptoms until 50% heading stage; T2 : NATIVO 300 SC at a dose of 350 ml.ha⁻¹, at the seedling (4 to 5 leaves), booting and complete heading stages; T3: NATIVO 300 SC at a dose of 350 ml.ha⁻¹, at seedling (4 to 5 leaves), maximum tillering and complete heading stages; T4: NATIVO 300 SC at a dose of 350 ml.ha⁻¹, at early tillering (2 to 3 stems visible), maximum tillering and complete heading stages). Data on disease severity of brown spot and agronomical parameters were recorded and statistically analysed using SAS 9.4 software. The results proved that treatments T1 and T4 proved more effective, with leaf severity indices of 2.0±0.98 and 3.52±0.78 versus 6.52±0.66 for the control, and yield gains of 67.69 and 60.09% respectively. In conclusion, application of NATIVO 300 SC fungicide at a dose of 350 ml/ha can be recommended at the early tillering, maximum tillering and complete heading stages of rice for effective management of brown spot.

Keywords: Chemical management; fungal disease; rice; phenological stages; environmental protection.

1. INTRODUCTION

Rice is a cereal that is consumed as a staple food in Côte d'Ivoire. In fact, the national average consumption of rice is estimated at 70 kg of milled rice per person, per year (Minader, 2019), while other staple foods are 17 kg per capita per year for wheat and 40 kg per capita per year for maize (ONDR, 2012). However, this cereal is insufficiently produced in Côte d'Ivoire. This gap in local production has been growing since the dissolution of SODERIZ at the end of the 1970s and will reach 50% by 2022. Faced with this situation of food insecurity, the government has been implementing a National Rice Development Strategy (SNDR) since 2012, with the short-term goal of achieving self-sufficiency by 2025 (ONDR, 2012). To achieve this, it will be necessary, among other things, to overcome the major biotic and abiotic constraints at the root of crop losses (Bouet et al., 2015). Among biotic constraints, fungal and viral diseases of rice reduce yields and increase production costs (Yang et al., 2020). Among fungal diseases, Brown spot of rice caused by a fungus, *Bipolaris oryzae* has been one of the most damaging to rice cultivation for about fifteen years (Bouet et al., 2015, Bouet et al., 2020, Bouet et al., 2022, Boka et al., 2018, Boka et al., 2020). Paddy losses due to brown spot have also been reported by several authors (Gupta et al., 2013, Balgude and Gaikwad, 2015, Sunil et al., 2017).

The occurrence of rice brown spot in almost all rice crops in Côte d'Ivoire threatens food security (Boka et al., 2020). In fact, this disease was the cause of the Bengal famine in India in 1942, which resulted in the deaths of more than two million people (Padmanabhan, 1973). Various phytosanitary products have been tested in several rice-producing countries to control rice brown spot (Bouet et al., 2020, Bouet et al., 2022, Boka et al., 2020, Gupta et al., 2013, Balgude and Gaikwad, 2015, Sunil et al., 2017, Karan et al., 2021). Some of these formulations have been shown to be effective against rice brown spot in Côte d'Ivoire. Although these products are effective, the financial implications of their use and their potential impact on the environment require intelligent approaches or recourse to other control methods. Moreover, the systematic use of chemical molecules exposes us to the risk of the emergence of resistant strains of pathogens (Gomathinayagam et al., 2011, Orou et al., 2024). The aim of this study was to rationalize the use of a fungicide effective against brown spot in a strict rainfed rice production system.

2. MATERIALS AND METHODS

2.1 Materials

The plant material used in this study is a widely cultivated local traditional cultivar "Gbékélé", with a crop duration of 105 days.

2.2 Methods

2.2.1 Study site

The trial was carried out in strict rainfed ecology at Biankouma (Côte d'Ivoire) between the months of July and November 2023. The area has been proved to be endemic for brown spot in recent decades (Bouet et al., 2020, Bouet et al., 2022). Geographically, the study site was referenced 07°40'36.2"N longitude, 007°36'25.9"W latitude at an altitude of 536 m.

2.2.2 Experimental design

A Fisher block design with three (03) replications were maintained. The fungicide NATIVO 300 SC (Tebuconazole 200 g/L +Trifloxystrobin 100 g/L) was used at a dose of 350 ml/ha for its efficacy against *Bipolaris oryzae*, the fungus responsible of rice brown spot according to the work of Bouet et al. (2022).

The treatment details of NATIVO 300 SC were as listed below:

- **T0**: no fungicide treatment;
- **T1**: NATIVO 300 SC at a dose of 350 ml.ha⁻¹, every 10 days interval from the initial appearance of brown spot symptoms until 50% heading stage;
- **T2**: NATIVO 300 SC at a dose of 350 ml.ha⁻¹, at the seedling (4 to 5 leaves), booting and complete heading stages;
- **T3**: NATIVO 300 SC at a dose of 350 ml.ha⁻¹, at seedling (4 to 5 leaves), maximum tillering and complete heading stages;
- **T4**: NATIVO 300 SC at a dose of 350 ml.ha⁻¹, at early tillering (2 to 3 stems visible), maximum tillering and complete heading stages.

A plot size of 12 m² were maintained.

2.2.3 Conduct of the trial

The trial was conducted by following all the package of practices from land preparation to weeding operations. Fungicide treatments were applied to these elementary plots according to the above fungicide application rates. The crop was harvested at maturity (paddy with 18% moisture content).

2.2.4 Data collection

Data on brown spot disease severity and agronomical data were recorded during the period of study. Disease data was scored according to IRRI's standard scoring scale (IRRI,

2014). Disease data was recorded at 10 days interval right from the initial appearance of the brown spot disease symptoms in the field. Agronomic parameters (fertile tillers, non-fertile tillers, plant height and 1000-grain weight) were recorded at maturity (paddy at 18% moisture). The paddy weights per elementary plot were determined after proper drying and winnowing. Data on Area under Brown Progression Curve (AUBSPC and grain yield were recorded:

- $AUBSPC = \sum_{n=i}^n (Y_{n+1} + Y_i) 0,5 (T_{n+1} + T_i)$
(where: Y_i = brown spot severity at i^{th} date; T_i = date on which the Y_i severity scoring was obtained; n = number of times where observations were made);
- $Yield\ gain\ (\%) = \frac{Yield\ Tf - Yield\ T0}{Yield\ T0} * 100$

Where Yield Tf = Yield of the fungicide treatment (T1, T2, T3 or T4) plots and Yield T0 = Yield of control plots (T0)

2.2.5 Statistical analysis

Data was analyzed using SAS 9.4 software (SAS, 1995). Brown spot severity indices and area under brown spot progression curve (AUBSPC) values, analysis of variances were performed using a general linear model (GLM), while for agronomic data, it was performed using one-factor ANOVA. The Newman-Keuls test (probability threshold $p=0.05$) was used for multiple comparisons.

3. RESULTS

3.1 Effect of Treatments on Brown Spot Severity at Different Stages of Crop Growth

At the early tillering stage (20 days after sowing), the first symptoms of the disease were observed on all plots before fungicide application, the severity index of brown spot was recorded for all treatments (Fig. 1).

At the maximum tillering stage, i.e. 45 days after sowing, the average severity index of brown spot in plots under treatment T0 was 2.33. For plots in treatment T1, the severity index of the disease was (1). For plots under treatments T2 and T3, the average severity index of brown spot was the same, i.e. (1.67). For plots under treatment T4, the average brown spot severity index was 1.33 (Fig. 1).




Fig. 1. Brown spot severity index progression according to rice development stage and the treatments

At the booting stage (75 days after sowing), the brown spot severity index for plots treated with T0 was 5.67. In plots receiving treatment T1, the disease severity index was 1.67. In plots receiving treatments T2 and T3, the same severity index of brown spot was observed with an average value of 3.00. For treatment T4, the average severity index of brown spot at the booting stage was 2.33 (Fig. 1).

At the heading stage (90 days after sowing), plots receiving treatment T0 showed an average severity index of 6.67 while it was 2.33 in the Treatment T1. But brown spot severity remains same in the Treatments T2 and T3 while it was 3.33 in T4 (Fig. 1).

3.2 Effect of NATIVO 300 SC on the Severity of Rice Brown Spot as per the Application Methods

According to the application modality of NATIVO 300 SC (Tebuconazole 200 g/L +Trifloxystrobin 100 g/L), the severity of brown spot recorded at rice maturity was 2.09 ± 0.98 for plots under treatment T1. For plots receiving treatment T2, the severity index of brown spot was 3.78 ± 1.45 . In plots receiving treatment T3, the average disease severity index was 3.52 ± 0.78 . In the plots under treatment T0 a high evolution of the disease was recorded, with an average severity index of 6.54 ± 0.66 (Table 1). The mean values of the area under the brown spot progression curve (AUBSPC) for the fungicide treatments ranged

from 93.57 ± 1.5 for treatment (T1) to 155.52 ± 0.48 for (T3), compared with a value of 270.5 ± 0.00 for control (T0).

Statistical analysis of severity indices and AUBSPC values recorded at the maturity stage showed a significant difference ($p=0.0001$) between fungicides treatments applied in this trial. Treatment (T1) was found to be more effective, with brown spot severity index of 2.09 ± 0.98 and AUBSPC mean value of 93.57 ± 1.5 . This treatment was followed by the fungicide treatment (T4), which showed a mean severity index of 3.52 ± 0.78 and AUBSPC mean value of 129.20 ± 0.83 . Treatments T2 and T3 had roughly the same effects, but weaker than those of treatments T1 and T4 (Table 1).

3.3 Effect of Treatments on Rice Agronomic Parameters

3.3.1 Effect of treatments on the number of total tillers

The results from the Table 2 showed that the number of total tillers varied from one treatment to another. The average number of total tillers recorded in fungicide-treated plots ranged from 7.04 ± 1.26 (T1) to 4.83 ± 1.80 (T4). The overall average number of total tillers was 6.5, with a coefficient of variation 19.42%. A significant difference ($p=0.0063$) was recorded between the fungicide treatments and the control, where the average number of total tillers was 7.52 ± 1.57 .

Table 1. Effect of NATIVO 300 SC (Tebuconazole 200 g/L +Trifloxystrobin 100 g/L) on Brown spot severity as per as application methods

Treatment	Leaf severity index	AUBSPC
T0	6.54±0.66a	270.07±0.00a
T1	2.09±0.98d	93.75±1.50d
T2	3.78±1.45b	155.27±1.27b
T3	3.85±0.60b	155.52±0.48b
T4	3.52±0.78c	129.20±0.83c
Overall mean	3.96±0.56	159.26±5.90
Probability (p) effect	0.0001 S	0.0001 S
CV(%)	18.14	0.612

AUBSPC: area under brown spot progression curve; CV (%): coefficient of variation; s: significant ($p < 0.05$). The values given in the table are the mean±standard deviation for each parameter. Means followed by the same letter are not significantly different at the probability threshold ($p = 0.05$) according to the Newman-Keuls test

3.3.2 Effect of treatments on the number of fertile tillers

The average number of fertile tillers observed in plots treated with NATIVO 300 SC fungicide (Tebuconazole 200 g/L + Trifloxystrobin 100 g/L) ranged from 6.5 (T1) to 3.70±1.72 (T4). In the control plots (T0), the average number of fertile tillers recorded was 6.59±1.49. The overall mean number of fertile tillers was 5.64±1.47, with a coefficient of variation 24.78%. There was a significant difference ($p=0.034$) between the different treatments applied (Table 2).

3.3.3 Effect of treatments on rice plant height

Average rice plant height ranged from 102.69±7.47 cm (T3) to 109.22±6.02 cm (T4) on plots treated with NATIVO 300 SC fungicide (Tebuconazole 200 g/L +Trifloxystrobin 100 g/L). For the control (T0), average height of rice plants was 102.80±5.28 cm. The overall mean of rice plants height was 106.09±6.90 cm, with a

coefficient of variation 7.84%. The statistical analysis results showed a significant difference ($p=0.0001$) between the treatments applied (Table 2).

3.3.4 Effect of treatments on yield and yield gain

Average paddy yields ranged from 0.63±0.16 t.ha⁻¹ for the control (T0) to 1.95±0.25 t.ha⁻¹ for the fungicide treatment (T1). The overall average yield for the trial was 1.43±0.48 t.ha⁻¹ with coefficient of variation 15.26%. Statistical analysis showed a significant difference ($p=0.0003$) between the yields of the fungicide-treated plots and the control plots (T0). No significant difference was observed between fungicides treatments for the yields. However, a yield gain of 67.69% was obtained with treatment (T1) followed by treatment (T4), which showed a yield gain of 60.37%. For treatments T2 and T3, the yield gains observed were 58.82% and 58% respectively (Table 3).

Table 2. Effect of NATIVO 300 SC (Tebuconazole 200 g/L +Trifloxystrobin 100 g/L) on agronomic parameters

Treatment	Total tillers	Fertile tillers	Heigh (cm)
T0	7.52±1.40a	6.59±1.49a	102.80±5.28b
T1	7.04±1.26ab	6.50±0.99a	107.30±5.07a
T2	6.66±1.94ab	6.03±1.87ab	108.58±8.26a
T3	6.48±1.12ab	5.37±1.26ab	102.69±7.47b
T4	4.83±1.80b	3.70±1.72b	109.22±6.02a
Overall mean	6.50±1.57	5.64±1.47	106.09±6.90
Probability (p) effect	0.0063 S	0.034 S	0.0001 S
C V (%)	19.42	24.78	7.84

CV (%): coefficient of variation; s: significant ($p < 0.05$). The values given in the table are the mean±standard deviation for each parameter. Means followed by the same letter or group of letters are not significantly different at the probability threshold ($p = 0.05$) according to the Newman-Keuls test

Table 3. Effect of NATIVO 300 SC (Tebuconazole 200 g/L +Trifloxystrobin 100 g/L) on Rice yields

Treatment	Yield (t.ha ⁻¹)	Yield gain (%)
T0	0.63±0.16b	-
T1	1.95±0.25a	67.69
T2	1.53±0.33a	58.82
T3	1.50±0.14a	58
T4	1.59±0.13a	60.37
Overall mean	1.43±0.48	
Probability (p)	0.0003	
effet	S	
CV(%)	15.26	

AUBSPC: area under brown spot progression curve; CV (%): coefficient of variation; s: significant (p=0.05). The values given in the table are the mean±standard deviation for each parameter. Means followed by the same letter or group of letters are not significantly different at the probability threshold (p= 0.05) according to the Newman & Keuls test




Fig. 2. Linear regression curve between yield and area under brown spot progression curve (AUBSPC) values

3.4 Regression between AUBSPC and Performance

A negative correlation was noted between yield and AUBSPC ($R^2 = 0.8447$). AUBSPC contributed 84.47% to yield variation, while other factors contributed for 15.53%. According to the linear regression equation for one unit of AUBSPC, yield is reduced by 0.0073 tonnes or 7.3 kg (Fig. 2).

4. DISCUSSION

Application of NATIVO 300 SC fungicide at 350 ml/ha at different phenological stages of the rice crop, induced a significant reduction of brown spot leaf severity. The experimental plots

received the Treatments T1 to T4. In the case of fungicide treatment (T1), application of NATIVO 300 SC at interval of ten days from the observation of the first symptoms of brown spot at the early tillering stage (20 days after sowing), reduced the brown spot leaf severity in rice to an average index of 2.09 and AUBSPC average value of 95.75, resulting in a yield gain of 67.69% compared with the control. Furthermore, when the fungicide treatments were applied at specific phenological stages, in the case of treatments T2, T3 and T4, a significant reduction of brown spot leaf severity was also observed. However, treatment T4 (application of the fungicide NATIVO 300 SC (Tebuconazole 200 g/L +Trifloxystrobin 100 g/L) at a dose of 350 ml/ha at the early tillering (2 to 3 stems visible),

maximum tillering and complete panicle heading stages) was more effective than treatments (T2 and T3). Treatment T4 reduced the disease severity index to 3.52, with AUBSPC average value of 129.20 and a yield gain of 60.37% compared to treatments T2 and T3, for which the observed leaf severity indices were 3.78 and 3.85, and AUBSPCs average values of 155.27 and 155.52 respectively. Average yield gains for these treatments were 58.82 and 58% respectively. For treatment T1, NATIVO fungicide was applied eight times from observation of the first symptoms to 50% heading. For the other treatments (T2, T3 and T4), the fungicide was applied three times.

The current results are in conformity with the work of Bouet et al. (2022) had shown that the fungicide NATIVO 300 SC (Tebuconazole 200 g/L +Trifloxystrobin 100 g/L) is effective at the dose of 350 ml/ha in reducing brown spot leaf severity. Treatment T1 was more effective, as it appears to prevent the complete development cycle of the fungal agent *Bipolaris oryzae* responsible of rice brown spot. In fact, the period during which this study was carried out was one of high humidity (75-80%), with an average temperature varying between 24° and 28°C, and therefore favorable to the development of the disease.

According to Datnoff & Lentini (1994), *Bipolaris oryzae* infection can occur between 20° and 26°C, and is favored by high relative humidity (86 to 100%) and the presence of free water on the plant surface. *Bipolaris oryzae* conidia germinate by producing a germ tube which penetrates the leaf through the epidermis or stomata. In the case of treatments (T2, T3 and T4) where the duration between two successive applications exceeded 25 days at least, the pathogen was able to establish itself adequately, as this duration allows *Bipolaris oryzae* to complete its life cycle. It has been found that rice leaf brown spot mainly attacks the late vegetative stages and becomes severe from the reproductive stage through to maturity, affecting photosynthetic activity where maximum translocation of sugars occurs from the leaves to the grains, which is responsible for the formation of light, shrivelled and straw-like grains and reduced yield (Viswanath et al., 2021). Also, during rice growth, fungus spores are formed on leaf spots and can subsequently infect other leaves and panicles (Nyvall et al., 1999). Severely infected leaves may die before maturity and these plants will produce light or empty grains (Agarwa et al., 1994, Kohls et al., 1987).

The low yield recorded in the control plots is therefore due to the reduction in the leaf area of rice plants. In the case of treatment T4, despite the low average number of fertile tillers compared with the other treatments, yield was high, as was yield gain. This could be explained by good grain filling in plots receiving this treatment.

Kohls et al. (1987) reported that yield losses specific to each growth stage and indicated that rice brown spot (*B. oryzae*) caused the greatest yield reductions when epidemics were triggered at the germination stage, and successively decreased when triggered late at the panicle initiation, heading and grain filling stages. Furthermore, Percich et al. (1997) demonstrated that the highest rates of infection by *B. oryzae* occurred at temperatures of 25 and 30°C and generally increased with periods of continuous humidity of 16, 18, 24 and 28 hours. Shafeeqa & Nandini, (2018) reported that fungicide treatments increased the number of tillers and the height of tillers compared with the absolute control. Furthermore, they also showed that NATIVO 75 WP fungicide at 70 ppm could extend the grain filling period by up to 6 days compared to the absolute control, maximizing grain yield. This could be due to ethylene production, which has the ability to decrease leaf and panicle senescence (Shafeeqa & Nandini, 2018). Balgude & Gaikwad, (2015) also showed that spraying a fungicide combination of trifloxystrobin 25% + tebuconazole 50% at a dose of 0.04% at 15-day intervals, starting the first spray immediately after disease onset, proved highly effective against brown spot, leaf blight, neck blight, knot blight, sheath rot and leaf scald, resulting in improved paddy grain yields. To ensure the safe application of trifloxystrobin and tebuconazole, Wang et al. (2024) conducted a study on their dissipation, terminal residues, distribution and associated risk factors. The results of their study indicated that trifloxystrobin and tebuconazole dissipate rapidly in rice, with an estimated half-life of between 4.1 and 7.7 days. The T4 treatment could therefore be recommended for rational use of the fungicide NATIVO 300 SC (a. i: Tebuconazole 200 g/L +Trifloxystrobin 100 g/L) at a dose of 350 ml/ha in the control of rice brown spot, in order to prevent the risk of environmental pollution and ensure good rice quality.

5. CONCLUSION

The current study was evaluated to know the effect of different application methods of the

fungicide NATIVO 300 SC (Tebuconazole 200 g/L +Trifloxystrobin 100 g/L) at a dose of 350 ml/ha on helminthosporiosis of rice. All fungicide treatments tested significantly reduced disease severity. However, treatments T1 and T4 found to be most effective, with yield gains of 67.69% and 60.09% respectively. Phenological stages that proved favorable for effective control of brown spot were early tillering, maximum tillering and heading. Application of the fungicide NATIVO 300 SC (Tebuconazole 200 g/L + Trifloxystrobin 100 g/L) i.e treatment T4 can be recommended to rice growing farmers at these phenological stages, as it not only reduces production costs but also reduce the dose of the fungicide at specific crop growth stage, thus protecting the environment. Looking ahead, a study needs to be carried out on the biochemical and organoleptic quality of rice produced under these conditions to confirm the absence of fungicide residues.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENTS

This study was carried out as part of the KAFACI-Sélection project "Development of new rainfed-lowland an irrigated varieties resistant to brown spot, RYWV and iron toxicity with good grain quality in Côte d'Ivoire" funded by the Republic of South Korea.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Agarwal, P. C., Mortensen, C. N., & Mathur, S. (1994). *Seed-borne diseases of rice and phytosanitary tests*. CIA, ADRAO. Sayce Publishing.
- Balgude, Y. S., & Gaikwad, A. P. (2015). Management of rice diseases with modern fungicides. *International Rice Symposium-IRS2015*, Nov 18–20, Hyderabad, 553.
- Boka, A., Bouet, A., Tiendrebeogo, A., Kassankogno, A. I., Ouedraogo, I., Ghislain, N. E., et al. (2018). Pathogenic variability of *Bipolaris oryzae* causing leaf spot disease of rice in West Africa. *International Journal of Phytopathology*, 7(3), 103–110.
- Boka, A., Bouet, A., Tonessia, C. D., Kouakou, M. B. K., & Denezon, O. D. (2020). Field evaluation of NATIVO 300 SC fungicide (Trifloxystrobin 100 g l-1 + Tebuconazole 200 g l-1) on rice brown spot (*Oryza sativa* L.). *Journal of Global Innovations in Agricultural Sciences*, 8(4), 177–182. <https://doi.org/10.22194/JGIASS/8.919>
- Bouet, A., Boka, A. K., Siapo, & Dogbo, O. D. (2022). Evaluation of the reactions of new rice genotypes to leaf spot (*Bipolaris oryzae*) in Côte d'Ivoire. *African Crop Science Journal*, 30(4), 405–414. <https://doi.org/10.4314/acsj.v30i4.8>
- Bouet, A., Gbedie, N. A., Boka, A., & Kouassi, N. (2015). Evaluation of promising rice varieties for resistance to some major biotic constraints and for their agronomic performances in Côte d'Ivoire. *International Journal of Biological and Chemical Sciences*, 9(4), 2041–2056.
- Bouet, A., Gueu, R. K., Boka, A., Noumouha, G. N. E., & Denezon, O. D. (2020). Field efficacy of ANTRACOL 70 WP, a fungicide based on Propineb 70%, under natural pressure of rice leaf spot caused by *Bipolaris oryzae*. *International Journal of Biological and Chemical Sciences*, 14(6), 2230–2239. <https://doi.org/10.4314/ijbcs.v14i6.24>
- Datnoff, L. E., & Lentini, R. S. (1994). Brown spot in Florida rice. *Plant Pathology Department, Institute of Food and Agricultural Sciences, University of Florida*, 128.
- Gomathinayagam, S., Balasubramanian, N., Shanmugaiah, V., Rekha, M., Manoharan, P. T., & Lalithakumari, D. (2011). Molecular characterization of carbendazim resistance of plant pathogen (*Bipolaris oryzae*). In N. Thajuddin & J. Trdine (Eds.), *Fungicides – Beneficial and Harmful Aspects* (pp. 197–218). Rijeka, Croatia: Sayce Publishing.
- Gupta, V., Shamas, N., Razdan, V. K., Sharma, B. C., Sharma, R., Kaur, K., Singh, I., John, D., & Kumar, A. (2013). Foliar application of fungicides for the management of brown spot disease in rice (*Oryza sativa* L.) caused by *Bipolaris oryzae*. *African Journal of Agricultural Research*, 8(25), 3303–3309. <https://doi.org/10.5897/AJAR2013.7182>
- IRRI. (2014). *Standard Evaluation System for Rice* (5th ed.). Manila, Philippines.

- Karan, R., Renganathan, P., & Balabaskar, P. (2021). In-vitro evaluation of selected fungicides against *Bipolaris oryzae* causing brown leaf spot of rice. *Research Journal of Agricultural Science*, 12(5), 1583–1584.
- Kohls, C. L., Percich, J. A., & Huot, C. M. (1987). Wild rice yield losses associated with growth-stage-specific fungal brown spot epidemics. *Plant Disease*, 71, 419–422.
- Minader. (2019). Aderiz highlights investment opportunities in the rice sector in Côte d'Ivoire. Press conference at the International Agricultural Show, Paris (France). Available at: http://www.agriculture.gouv.ci/uploads/Au_SIA_2019_I%E2%80%99ADERIZ_met_en_exergue_les_opportunit%C3%A9s_d%E2%80%99investissement_dans_la_fili%C3%A8re_riz
- Nyvall, R. F., & Percich, J. A. (1999). Development of fungal brown spot and spot blotch on cultivated wild rice in Minnesota. *Plant Disease*, 83, 936–938.
- ONDR (National Office for the Development of Rice Cultivation). (2012). *Revised National Strategy for the Development of the Rice Sector in Côte d'Ivoire (SNDR) 2012-2020*. Abidjan: MINAGRI.
- Orou, R. K., Yapo, W. T., Tanoh, G. A., Onetie, O. Z., & Orou, M. F. O. (2024). Study of farmer poisonings due to agricultural pesticides in the Agboville department, Ivory Coast. *International Journal of Biological and Chemical Sciences*, 18(2), 405–413. <https://doi.org/10.4314/ijbcs.v18i2.7>
- Padmanabhan, S. Y. (1973). The great Bengal famine. *Annual Review of Phytopathology*, 11, 11–26.
- Percich, J. A., Nyvall, R. F., Malvick, D. K., & Kohls, C. L. (1997). Interaction of temperature and moisture on infection of wild rice by *Bipolaris oryzae* in the growth chamber. *Plant Disease*, 81, 1193–1195.
- SAS. (1995). *Guide to the use of PC-SAS Version 6.04 for DOS for Statistical Analysis*. Cary, North Carolina: SAS Institute.
- Shafeeqa, T., & Nandini, K. (2018). Effect of eco-friendly chemicals on morphological and phenological characters of rice variety Uma under UV stress. *Agricultural Science Digest*, 38(3), 172–177.
- Sunil, S., Laxman, A., Binaya, P., Jharana, P., Pratibha, S., & Yogendra, S. S. (2017). Field experiment to evaluate the efficacy of different doses of chemical fungicides against rice brown leaf spot disease caused by *Bipolaris oryzae* L. at Paklihawa, Rupandehi, Nepal. *World Journal of Agricultural Research*, 5(3), 162–168. <https://doi.org/10.12691/wjar-5-3-6>
- Viswanath, H. S., Singh, R., Singh, G., Mishra, P., Shahi, U. P., Singh, D. V., et al. (2021). Impact of agro-met conditions and crop growth stages on the progression of brown spot disease in basmati rice. *International Journal of Environment and Climate Change*, 11(7), 59–67. <https://doi.org/10.9734/IJECC/2021/v11i730440>
- Wang, S., Liu, Y., & Zhu, M. (2024). Safe application of 75% trifloxystrobin–tebuconazole as water-dispersible granules in paddy based on residue and dietary risk assessment. *Molecules*, 29, 163. Available at: <https://doi.org/10.3390/molecules29010163>
- Yang, F., Zhang, J., Zhang, H., Ji, G., Zeng, L., Li, Y., Yu, C., Fernando, W. G. D., & Chen, W. (2020). Bacterial blight induced shifts in endophytic microbiome of rice leaves and the enrichment of specific bacterial strains with pathogen antagonism. *Frontiers in Plant Science*, 11, 963. <https://doi.org/10.3389/fpls.2020.00963>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/125337>