



Avocado Peel Polyphenolic-Rich Extract Attenuates Inflammatory Responses in Alloxan-Induced Diabetic Male Wistar Rats

George O. Eidangbe ^a and Olarewaju M. Oluba ^{b*}

^a Department of Medical Biochemistry, College of Medicine, Ambrose Alli University, Ekpoma, Edo State, Nigeria.

^b Department of Medical Biochemistry, David Umahi Federal University of Health Sciences, Uburu, Ebonyi State, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajarr/2024/v18i12841>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/127435>

Original Research Article

Received: 14/10/2024
Accepted: 18/12/2024
Published: 21/12/2024

ABSTRACT

Aim: This study sought to assess the effects of polyphenolic-rich extract of the peel of *Persea americana* (avocado) (APPE) on inflammatory responses in alloxan-induced diabetic rats. **Methodology:** A total of 35 alloxan-induced diabetic male rats were randomly divided into five groups (n = 7) and designated: Diabetic Control (DC), and groups receiving 25 (APPE₂₅), 50 (APPE₅₀), 100 (APPE₁₀₀), and 200 (APPE₂₀₀) mg/kg of APPE. In addition, a normal control (NC) treated with distilled water was maintained. Treatment was administered orally once daily for 21 days.

*Corresponding author: Email: olubamichael2023@gmail.com, olubamo@dufuhs.edu.ng;

Cite as: Eidangbe, George O., and Olarewaju M. Oluba. 2024. "Avocado Peel Polyphenolic-Rich Extract Attenuates Inflammatory Responses in Alloxan-Induced Diabetic Male Wistar Rats". *Asian Journal of Advanced Research and Reports* 18 (12):436-49. <https://doi.org/10.9734/ajarr/2024/v18i12841>.

Results: After 21 days, the APPE-treated rats, particularly at 200 mg/kg, exhibited significant ($p < 0.05$) improvements in both body weight and blood glucose levels compared with DC group. Significant ($p < 0.05$) reductions in glycated hemoglobin (GLH), interleukins-1 (IL-1), and (interleukin-6 IL-6), and tumor necrosis factor- α (TNF- α) levels were observed in the APPE-treated rats compared with DC. In contrast, the APPE-treated rats exhibited significant ($p < 0.05$) improvement in insulin and interleukin-10 levels compared with DC rats. Furthermore, DC and APPE₂₅ and APPE₅₀ showed varying degrees of interfacial hepatitis and glomerulosclerosis. Nevertheless, treatment with APPE₂₀₀ exhibited protection against the detrimental impact of diabetes mellitus on the liver and kidney in alloxan-induced diabetic rats.

Conclusion: Avocado peel polyphenolic-rich extract at 200 mg/kg body weight effectively inhibited low-grade inflammation induced by oxidative stress in the liver and kidneys of rats exposed to alloxan.

Keywords: Avocado peel extract; polyphenolic extract; cytokines; inflammation; diabetic complications.

1. INTRODUCTION

Persistent high blood sugar level is a major predictive determinant in the development of diabetes and its associated complications. Metal-ion-catalysed glucose oxidation potentiates the production of free radicals including hydrogen peroxide and ketoaldehydes. Ketoaldehyde in turn enhances the generation of superoxide, which is spontaneously converted to H₂O₂ and subsequently into the remarkably powerful hydroxyl radical (Pamplona, 2008). The overwhelming effects of the continuous production of these free radicals in diabetes on biological antioxidants capacity leads to the emergence of oxidative stress-induced complications in individuals with diabetes (Wu et al., 2011).

Consequently, there exists a correlation between oxidative stress and the emergence of macrovascular problems in individuals with type 2 diabetes (Bigagli and Lodovici, 2019). One of the primary mechanisms implicated in the non-enzymatic glycosylation of hemoglobin is the autooxidation of glucose. Hemoglobin that has been glycosylated is a powerful generator of free radicals. Therefore, there exists a causal relationship between diabetic complications, hemoglobin glycosylation and oxidative stress. Nevertheless, the aforementioned processes may be mitigated subsequent to the administration of antioxidants.

The tropical fruit avocado pear (*Persea americana* Mill. Lauraceae), sometimes referred to as alligator pear or butter fruit, possesses flesh that can be consumed. The waste generated during the production of avocado pear ranges from 35% to 50%, with the peel accounting for around 20% to 25% of this waste

(Figuroa et al., 2018). Nevertheless, it has been documented that avocado peel possesses a substantial quantity of phenolic chemicals. The phenolic content of avocado peel waste has been found to be much higher in comparison to avocado pulp (Vinha et al., 2020). Research has demonstrated that the phenolic extract derived from avocado pear exhibits a wide range of pharmacological properties, encompassing anti-diabetic (Oguntibeju, 2019) anti-inflammatory, and antioxidant effects (Alkhalaf et al., 2019). Avocado peel waste is regarded as a promising and cost-effective biomass for extracting phenolic compounds from agricultural waste, due to its numerous health benefits. Therefore, the utilization of avocado peel waste holds significant potential for enhancing the value of the avocado processing industry and effectively mitigating the environmental consequences associated with avocado waste (Oluba et al., 2019). Recent reports indicate that specific foods and plant extracts have the potential to hinder the activity of α -amylase. The inhibition observed in these foods and plant extracts can be attributed to the presence of polyphenols, as documented (Colombo et al., 2019). Considering the potential existence of a causal association between oxidative stress induced by hyperglycemia and the glycosylation of hemoglobin, as well as the subsequent induction of inflammatory cytokines in individuals with diabetes (National Research Council, 2010, Percie du Sert et al., 2020). We predicted that the potential inhibition of carbohydrate hydrolyzing enzymes and/or the probable suppression of glucose oxidation via antioxidant mechanisms by avocado peel polyphenolic-rich extract could offer a viable method in the prevention and management of diabetes complications.

2. MATERIALS AND METHODS

2.1 Chemicals and Reagents

Chemicals and reagents used in this study were AnalaR grades and were products of Merck Life Science UK Ltd. (Gillingham, United Kingdom) except as stated otherwise.

2.2 Plant Materials

Matured *Persea americana* (avocado pear) fruits were collected from a vegetation area near Omu-Aran, Kwara State, Nigeria (8°8'18" N; 5°6'9" E) after obtaining permission from the farm owner to collect the fruits between July 2020 and August 2020. Mr. Bolu of the University of Ilorin's Department of Plant Science, Ilorin, Nigeria, where the voucher specimen (UILH/011/1803) was deposited, identified and authenticated the plant sample.

2.3 Animals

Male Wistar rats weighing 180–200 g procured from the animal holding unit of the Department of Biochemistry, Landmark University, Omu-Aran, Nigeria, were used for the study. The animals were housed in standard rat cages with seven animals per cage with a 12 h light/dark cycle and fed *ad libitum* with unrestricted access to water. Prior authorization for animal handling was sought from the Landmark University, Omu-Aran, Nigeria Research and Ethics Committee (approval number LMU/EC/098/2019).

2.4 Microwave Assisted Polyphenol Extraction

The modified microwave-assisted method of Simi et al., 2016 was used to extract total polyphenols with little modification. Briefly, in a 500-mL Erlenmeyer flask containing 250 mL of ethanol (50%), powdered avocado pear peel (50 g) was placed and put in a BP090 microwave oven (Microwave Research & Applications, Inc., Illinois, USA). A vertical condenser was securely connected to the flask. For 5 minutes, the extraction process was carried out at 300 W, and the liquid extract was separated from the residue using vacuum filtration and vacuum evaporated at 40 °C to obtain the concentrated polyphenolic-rich extract.

2.5 Determination of Total Polyphenol

The total polyphenol content of avocado peel polyphenolic extract was calculated using Ozcan

et al. colorimetric Folin-Ciocalteu method (Özcan et al., 2020).

2.6 GC-MS Analysis

The preparation of the avocado peel polyphenolic extract was carried out according to the Oluba et al. method (Oluba et al., 2020). A weighed amount of the avocado peel polyphenolic extract was dissolved in GC-grade methanol (1:5 w/v). The mixture was shaken in an orbital shaker for 48 hours at 252 °C. Thereafter, the mixture was filtered through a muslin cloth to obtain the crude extract, which was concentrated using a rotary evaporator and stored at -4 °C until required for further analysis. The methanol extract was then subjected to GC-MS analysis in order to identify its constituent bioactive metabolites. The detection was carried out with an ionisation energy of 70 eV and 60 kPa using helium as the carrier gas. The oven temperature was initially set at 100 °C for 2 min with a ramp rate of 4 °C per min to 225 °C and then 1 °C per min to 245 °C, followed by 40 °C to 280 °C with a 30 min hold. Thereafter, a 2 µL were injected at 280 °C with a split ratio of 1:50. The detected spectra were identified from the NIST database using the ChemStation software.

2.7 Induction of Diabetes

A single intraperitoneal administration of 150 mg/kg alloxan monohydrate in normal saline was administered to the rats (Macdonald et al., 2017). Seventy-two hours after alloxan administration, rats were tested for fasting blood glucose levels from tail vein blood using a One-Touch Ultramini blood glucose meter (LifeScan, Milpitas, USA). Animals with 200 mg/dL and above fasting blood glucose were considered diabetic and included in the study.

2.8 Experimental Design

The experimental design consisted of 35 alloxan-induced diabetic rats and a separate set of 7 non-diabetic rats. The animals were grouped as follows:

Normal control (NC): Normal rats ($n = 7$) treated with distilled water,

Diabetic control (DC): Alloxan-diabetic ($n = 7$) treated with distilled water,

APPE₂₅: Alloxan-diabetic rats ($n = 7$) treated with APPE (25 mg/kg bw),

APPE₅₀: Alloxan-diabetic rats ($n = 7$) treated with APPE (50 mg/kg *bw*),

APPE₁₀₀: Alloxan-diabetic rats ($n = 7$) treated with APPE (100 mg/kg *bw*),

APPE₂₀₀: Alloxan-diabetic rats ($n = 7$) treated with APPE (200 mg/kg *bw*).

Treatment was given orally once daily for 21 consecutive days, after which they were euthanized with the inhalation of isoflurane after which blood collected by cardiac puncture. Body weight and fasting blood glucose level were taken every three days. After the 21st-day treatment, rats in each group were sacrificed under anesthesia, and blood was collected into clean, sterile tubes while intestine, liver, heart, and kidney samples were quickly excised, blotted with tissue paper, freed of fats, and stored in formalin.

2.9 Blood Sugar Determination

Blood was taken from cut tip of the respective rat tail and fasting blood sugar level was estimated using One Touch Ultramini glucometer.

2.10 Determination of Glycated Hemoglobin

Hemoglobin glycosylation level was evaluated following Nayak and Pattabiraman, 1981. Method Briefly, whole blood sample was combined with a lysing reagent that consisted of a detergent and borate ions. This ensured the removal of the unstable Schiff's base during the process of hemolysis. The hemolysate was thoroughly mixed for a minimum of 5 minutes with a cation exchange resin that has a low affinity for binding. At this moment, HbA₀, which is the form of hemoglobin that is not glycated, is attached to the resin. A specialized resin separator was employed to extract the resin from the supernatant fluid, which included the HbA₁ (glycated HbA). The percentage of glycated hemoglobin (GLH) in relation to total hemoglobin was calculated by measuring the absorbance of the glycohemoglobin and total hemoglobin at 415 nm and comparing it to a reference glycohemoglobin preparation used in the test.

2.11 Biochemical Analysis

Whole blood was allowed to stand at room temperature (25 ± 2 °C) for 30 min before being centrifuged at 1500 *xg* for ten minutes in a

refrigerated centrifuge, and serum was carefully pipetted into dry, clean, sterile bottles and stored at -4 °C for further biochemical analyses. Serum interleukins (IL-1 β , IL-6, and IL-10) and tumour necrosis factor- α (TNF- α) levels, were determined using their respective diagnostic kits (R&D System, Minneapolis, USA) in strict accordance with the manufacturer's instructions.

2.12 Histological Evaluation

A small portion of the liver sample was cut and fixed in 10% neutral buffered formalin. For light microscopy, 5 μ m thin sections were cut with a rotary microtome and stained with haematoxylin and eosin. The slides were viewed with an appropriate objective to check for the architecture of the tissues. The images were shown in photomicrographs, and the histology was examined.

2.13 Statistical Analysis

Data were expressed as means \pm SEM. Statistical analysis and graphical representation were carried out using GraphPad Prism software (version 8.0). The mean comparison was done by one-way analysis of variance (ANOVA), followed by Least Significant Difference (LSD) test. *P* values less than 0.05 were considered significant.

3. RESULTS AND DISCUSSION

3.1 Phytochemical Composition

The total phenolic content of avocado peel polyphenolic extract was 127.1 mg garlic acid equivalent⁻¹ sample. The GC-MS fingerprint of avocado peel polyphenolic extract is shown in Fig. 1. According to the GC-MS profile, the following phytochemicals were identified, butanoic acid, 4-chloro-, 1,1a,1b,4,4a,5,7a,7b,8,9-decahydro-4a,7b-dihydroxy-3-(hydroxy methyl)-1,1,6,8 -tetramethyl-5-oxo-9aH-cyclo propa[3,4]benz[1,2-e]azulene-9,9a-diy ester, [1a-(1 α ,1b β ,4a β ,7a α ,7b α ,8 α ,9 β ,9a α)] -; Card-20(22)-enolide,3-[(2,6-dideoxy-4-O- β -D-glucopy ranosyl-3-O-methyl- β -D-ribo-hexopy ranosyl) oxy]-5,14-dihydroxy- 19-oxo-, (3 β , 5 β)-;2,5-Dichloro-3,6-bis-(1-methyl-1H-benzoimidazol-2-yl)-terephthalic acid, dimethyl ester; 5H-Cyclopropa[3,4]benz[1,2-e]azulen-5-one, 3,9,9a-tris(acetyloxy)- 3-[(acetyloxy)methyl]- 2-chloro- 1,1a,1b,2,3,4,4a,7a,7b,8,9,9a-dodecahy dro-4a,7b-dihydroxy- 1,1,6,8-tetramethyl-, 1aR-

(1 α ,1 β ,2 α ,3 β ,4 α ,7 α ,7 β , 8 α ,9 β ,9 α)-; Bufa-14,16,20,22-tetraenolide, 3-(acetyloxy)-, (3 β ,5 β)-; Azafrin; Cholestane, 3,5-dichloro-6-nitro-, (3 β ,5 α ,6 β)-; and Chlordiazepoxide (Table 1). These identified molecules are either are either steroids, polyphenolics, terpenes or fatty esters, etc. The results show that quercetin was the major phenolic compound present in avocado pear peel. This observation is consistent with the report of Rosero et al.,2019 From previous studies, quercetin has been proven to inhibit tissue oxidative damage via its augmenting effect on oxidative status. Moreover, it has been established that it enhances the regeneration of pancreatic islets (Abdelkader et al., 2020). In addition, plant phenolics have been indicated to inhibit the action of α -amylase in the digestion of carbohydrates in diabetes, while others have been proven to lower the risk of complications arising from type 2 diabetes (Yang and Kang, 2018). Flavonoids, phenolic acids, and tannins have been shown to inhibit the actions of the two main enzymes involved in the breakdown of dietary carbohydrates, α -glucosidase and α -amylase (Abdelkader et al., 2020).

3.2 Effect on Body Weight





At the end of the treatment period, a significant decrease in body weight (-32.3%) was observed in alloxan-induced diabetic rats (DC) in comparison with that of the normal control group. Interestingly, treatment of diabetic animals with avocado peel polyphenolic-rich extract at 50, 100, and 200 mg/kg body weight (i.e. APPE₅₀, APPE₁₀₀, and APPE₂₀₀) led to a significant ($P = .05$) increased in body weight compared to





diabetic control (DC). APPE₂₀₀ gave the most significant ($P = .05$) increase in body weight, followed by APPE₅₀ and APPE₁₀₀, which gave a similar effect in body weight at the end of the treatment period. No significant difference in body weight was observed between APPE₁₀₀ and APPE₅₀ at the end of the treatment period (Fig. 2). Alloxan selectively destroys pancreatic islet Langerhans -cells, resulting in a massive reduction in insulin production, hyperglycemia, and a subsequent derangement in energy metabolism (Rodrigues et al., 2018, Yang et al., 2021) as vividly observed in this study. In compliance with previous studies, the present study showed that alloxan intoxication in rats led to reduced body weight while enhancing food consumption. Though data on food intake is not reported in this study, alloxan-induced rats were observed to consume more food than non-diabetic control rats. The usual decrease in body weight in a diabetic state could be attributed to reduced cellular uptake and utilisation of blood glucose, which also influence the rate of lipid dispensation and gluconeogenesis (Yang and Kang, 2018). The increase in body weight in alloxan-induced diabetic rats administered avocado peel polyphenolic-rich extract could have been a resultant anabolic effect of its phenolic phytochemicals, which are capable of abrogating the physiological consequences of alloxan and hyperglycemia-induced oxidative stress through their antioxidant activity. Dietary plant polyphenols have been shown to improve hyperglycemia, dyslipidemia, and insulin resistance by regulating carbohydrate and lipid metabolism (Shahwan et al., 2022).



Fig. 1. GC-MS fingerprint of avocado peel polyphenolic extract

Table 1. Phytochemical components of avocado pear peel polyphenolic extract obtained using GC-MS

s/n	Retention time (min)	Peak height	% of total	Molecular formula	Exact mass	Name	Structure
1	5.956	725385	1.203	C ₂₈ H ₃₈ Cl ₂ O ₈	572.19	Butanoic acid, 4-chloro-, 1,1a,1b,4, 4a,5,7a, 7b,8,9-decahydro-4a,7b-dihydroxy-3-(hydroxymethyl)-1,1,6,8-tetramethyl-5-oxo-9aH-cyclopropa[3,4]benz[1,2-e]azulene-9,9a-diylester,[1a α ,1b β ,4a β ,7a α ,7b α ,8 α ,9 β ,9a α]-	
2	6.881	858344	1.628	C ₃₆ H ₅₄ O ₁₄	710.35	Card-20(22)-enolide,3-[(2,6-dideoxy-4-O- β -D-glucopyranosyl-3-O-methyl- β -D-ribo-hexopyranosyl)oxy]-5,14 -dihydroxy-19-oxo-, (3 β ,5 β)-	
3	7.288	1578473	1.617	C ₂₆ H ₂₀ Cl ₂ N ₄ O ₄	522.09	2,5-Dichloro-3,6-bis-(1-methyl-1H-benzoimidazol-2-yl)-terephthalic acid, dimethyl ester	
4	8.289	1974119	1.737	C ₂₈ H ₃₇ ClO ₁₁	584.20	5H-Cyclopropa[3,4]benz[1,2-e]azulen-5-one,3,9,9a-tris(acetyloxy)-3-[(acetyloxy)methyl]-2-chloro-1,1a,1b,2,3,4,4a,7a,7b,8,9,9a-dodecahydro-4a,7b-dihydroxy-1,1,6,8-tetramethyl-, [1a α ,1b β ,2 α ,3 β ,4a β ,7a α ,7b α ,8 α ,9 β ,9a α]-	

s/n	Retention time (min)	Peak height	% of total	Molecular formula	Exact mass	Name	Structure
5	9.506	26.83217	3.637	C ₂₆ H ₃₂ O ₄	408.23	Bufa-14,16,20,22-tetraenolide, 3-(acetyloxy)-, (3β,5β)-	
6	11.782	46.82903	2.664	C ₂₇ H ₃₈ O ₄	426.28	Azafrin	
7	12.561	8473565	6.024	C ₂₇ H ₄₅ Cl ₂ NO ₂	485.28	Cholestane, 3,5-dichloro-6-nitro-, (3β,5α,6β)-	
8	12.882	3424344	1.808	C ₁₆ H ₁₄ ClN ₃ O	299.08	Chlordiazepoxide	




Fig. 2. Effect of avocado pear (*Persea americana*) polyphenolic peel extract (APPE) on body weights in alloxan-induced diabetic rats

Results are means \pm SEM of five replicates. Note: NC, normal control; DC, diabetic control; 25 mg/kg, 50 mg/kg, 100 mg/kg and 200 mg/kg, alloxan-induced diabetic rats treated with 25, 50, 100 and 200 mg/kg APPE, respectively

3.3 Effects on Blood Glucose, insulin and Glycated Hemoglobin Concentrations

In comparison to normal control, diabetic control had a significant ($P = .05$) increase in serum glucose level (241.4%). APPE₅₀, APPE₁₀₀, and APPE₂₀₀-treated diabetic rats had significantly lower serum glucose concentrations ($P = .05$) when compared to DC. At the end of the treatment period, APPE₂₀₀ provided the most significant ($P = .05$) reduction in blood glucose, followed by APPE₁₀₀ and APPE₅₀, which provided a similar reduction in glucose level. At the end of the treatment period, there was no significant ($P = .05$) difference in blood glucose levels between APPE₂₀₀ and NC (Fig. 3a). When comparing DC to NC, there was a significant drop in serum insulin concentration (69.1%). However, APPE-treated rats showed a dose-dependently significant ($P = .05$) increase in serum insulin levels when compared to control rats (Fig. 3b). When comparing DC to NC, there was a significant ($P = .05$) increase (96.4%) in GLH concentration. However, APPE-treated rats had a dose-dependently significant ($P = .05$) decrease in GLH concentration when compared to DC (Fig. 3c). The hallmark of diabetic management is the restoration of blood glucose levels to normal. As observed in the present study, avocado peel polyphenolic-rich extract at

200 mg/kg body weight proved to be an effective hypoglycemic agent in decreasing blood glucose concentrations to their normal value in alloxan-induced diabetic rats. As a result, avocado peel polyphenolics may be responsible for producing or amplifying a beneficial role in carbohydrate utilisation in alloxan-induced diabetic rats.

The results in this study showed a significantly higher level of GLH in diabetic but untreated rats (DC) compared to both the normal control and diabetic groups treated with avocado peel polyphenolic-rich extract. This shows that haemoglobin becomes glycosylated in a condition of high blood glucose concentration. The reduction in the level of GLH in the avocado peel polyphenolic-rich extract-treated diabetic rats clearly suggests that avocado peel polyphenolic treatment reduces protein glycosylation in diabetic rats. The inhibitory action of avocado peel polyphenolics on protein glycosylation could be attributed to the antioxidative effects of polyphenolics through Millard reaction. Several antioxidants, including ascorbic acid and β -carotene, have been shown *in vivo* and *in vitro* to inhibit protein glycosylation (Berryman et al., 2005, Zafar et al., 2012). Dietary antioxidants have also been reported to be capable of scavenging free

radicals produced by protein glycosylation (Perez et al., 2012).

3.4 Effects of Serum Cytokine Concentration

A significant ($P = .05$) increase (96.4%) in serum IL-1 β concentration was observed in DC compared to NC. However, a dose-dependently significant ($P = .05$) decrease in serum IL-1 β concentration was observed in the APPE-treated animals compared to DC (Fig. 4a). A significant ($P = .05$) increase (96.4%) in serum IL-6 concentration was observed in DC compared to NC. In contrast, a dose-dependently significant ($P = .05$) decrease in serum IL-6 concentration was observed in the APPE-treated rats up to 100 mg/kg compared to DC. IL-6 level was not significantly different in APPE₁₀₀ and APPE₂₀₀ (Fig. 4b). When comparing DC to NC, there was a significant ($P = .05$) increase (96.4%) in serum TNF- α concentration. However, APPE-treated rats had a dose-dependently significant ($P = .05$) decrease in serum TNF- α concentration when compared to DC. The difference in TNF- α concentrations between APPE₂₅ and APPE₅₀ was not statistically significant (Fig. 4c). A marked drop in serum IL-10 concentration was

observed in DC compared to NC. However, a dose-dependently significant ($P = .05$) decrease in serum IL-10 concentration was observed in APPE₅₀ to APPE₂₀₀ compared to DC. The observed difference in IL-10 concentrations in APPE₂₅ and APPE₅₀ was not significant (Fig. 4d). Oxidative stress has been postulated to be capable of activating several pathological changes in almost every type of kidney cell, including endothelial cells, mesangial cells, tubular cells, etc. This process in turn results in the development of diabetic neuropathy. It has also been pointed out that fibrosis, which appears to be the most prominent feature of diabetic neuropathy, is closely linked to inflammation (Wada and Makino, 2013). In accordance with previous postulations, the serum level of pro-inflammatory cytokines was observed to be elevated in alloxan-induced diabetic rats in this study (Domingueti et al., 2016, Domingueti et al., 2018). This study discovered that serum pro-inflammatory molecules like IL-1 β , IL-6, and TNF- α were upregulated in alloxan-induced diabetic rats. On the contrary, there was a corresponding reduction in serum concentrations of the anti-inflammatory cytokine IL-10.




Fig. 3. Effect of avocado pear (*Persea americana*) peel polyphenolic extract (APPE) on (a) blood glucose, (b) insulin and (c) glycated hemoglobin concentrations in alloxan-induced diabetic rats. Results are means \pm SEM of five replicates

Note: NC, normal control; DC, diabetic control; 25 mg/kg, 50 mg/kg, 100 mg/kg and 200 mg/kg, alloxan-induced diabetic rats treated with 25, 50, 100 and 200 mg/kg APPE, respectively. Bars carrying different Greek alphabets are significant ($P = .05$)




Fig. 4. Effect of avocado pear (*Persea americana*) peel polyphenolic extract (APPE) on serum (a) interleukins-1 β , (b) interleukin-6, (c) tumour necrosis factor- α , and (d) interleukin-10 concentrations in alloxan-induced diabetic rats

Results are means \pm SEM of five replicates. Note: NC, normal control; DC, diabetic control; 25 mg/kg, 50 mg/kg, 100 mg/kg and 200 mg/kg, alloxan-induced diabetic rats treated with 25, 50, 100 and 200 mg/kg APPE, respectively. Bars carrying different Greek alphabets are significant ($P = .05$)




Fig. 5. Haematoxylin and eosin stained photomicrographs of the (a) intestine, (b) heart, (c) liver, and (d) kidney

Note: NC, normal control, non-diabetic rats administered distilled water; DC, diabetic control, alloxan-induced diabetic rats administered distilled water; 25, 50, 100, and 200, alloxan-induced diabetic rats administered 25, 50, 100, and 200 mg/kg body weight, respectively avocado pear peel polyphenolic extract. Magnification X400

Recently, the concentrations of these pro-inflammatory molecules were observed to increase and correlate positively with the progression of diabetic nephropathy (Yaribeygi et al., 2019). These observations give an indication of a possible direct link between these inflammatory markers and glomerular damage. According to Akdis et al., 2016. IL-10, also known as "cytokine synthesis inhibitory factor," is produced by T-cells (CD4⁺), which play a significant role in adaptive immunity, and inhibits the synthesis of pro-inflammatory cytokines such as IL-4 and IL-5 (from Th2 cells) as well as the production of IL-2 and interferon- γ (IFN- γ) by Th1 cells. Thus, IL-10's anti-inflammatory role includes inhibitory actions on both adaptive and innate immune cells (Rouhi et al., 2020). It is believed that the synthesis and secretion of these fibrogenic cytokines in the local microenvironment may be responsible for the observed damage in renal architecture (Sindhughosa and Pranamartha, 2017). Furthermore, the ability of these molecules to recruit circulating red blood cells and direct their migration into kidney tissue has been identified as a possible risk factor for diabetic nephropathy.

3.5 Effect on Cellular Architecture

The cytological architecture of the intestine and heart across the different groups were observed to be normal at the end of the treatment period. However, photomicrographs of the liver and kidney showed interface hepatitis (a type of liver inflammation and necrosis that occurs at the junction of the portal tract and the hepatic parenchyma), and glomerulosclerosis, defined by the scarring of the glomeruli due to the interaction of infiltrating inflammatory cells with the glomerular cells, respectively, in alloxan-induced but untreated diabetic rats. Treatment with avocado peel polyphenolic-rich extract at varying concentrations showed a dose-dependent improvement in the liver and kidney. Avocado peel polyphenolic-rich extract at 200 mg/kg restored the liver and kidney architecture to normal at the end of the treatment period (Fig. 5). The histological examinations of the intestine, heart, liver, and kidney revealed intact architecture in all the tissues across the various treatment groups except in the kidney, where varying degrees of interface hepatitis, and glomerulosclerosis, were observed in the alloxan-induced diabetic rats. However, avocado peel polyphenolic-rich extract at 100 and 200 mg/kg body weight abrogated these abnormalities and restored the kidney

architecture. These findings further reinforced the claim that diabetic nephropathy is the most prominent feature of diabetic complications.

Based on the findings of this study, it is hypothesised that the induction of diabetes in rats using alloxan, resulting in sustained hyperglycemia due to pancreatic beta cell destruction and the subsequent non-enzymatic glycosylation of hemoglobin, could have been responsible for the stimulation, recruitment, and release of pro-inflammatory cytokines (IL-1 β , IL-6 and TNF- α) thus potentiating kidney damage. IL-1 β , IL-6 and TNF- α have been shown to activate the **nuclear factor kappa-B (NF- κ B) pathway, which regulates inflammatory, immune function, and stress responses**, thereby mediating the phosphorylation of serine molecules in the insulin receptor substrate (Ajiboye et al., 2018). Insulin resistance is thought to be exacerbated by phosphorylation of the insulin receptor substrate (Gratas-Delamarche et al., 2014). However, avocado peel polyphenolic extract significantly reduced these effects, particularly at 200 mg/kg body weight, probably via its inhibitory effects on carbohydrate hydrolyzing enzymes (Yang and Kang, 2018) as well as its antioxidant activity in preventing the oxidation of glucose and the generation of reactive glycated proteins.

4. CONCLUSION

This study found that a polyphenol-rich extract of *Persea americana* peel reduced the sustained hyperglycemia caused by alloxan administration in rats and its attendant oxidative stress-induced-low-grade inflammation by decreasing serum levels of glycated hemoglobin and pro-inflammatory cytokines such as IL-1 β , IL-6, and TNF- α and increasing serum concentrations of serum IL-10, an anti-inflammatory cytokine. The restoration in renal cellular architecture observed in diabetic rats treated with the extract provided further justification for this claim.

DISCLAIMER

This paper is an extended version of a preprint document of the same author.

The preprint document is available in this link:

<https://www.researchsquare.com/article/rs-910440/v1>

<https://assets.researchsquare.com/files/rs-910440/v1/356a4f74-c7e6-4452-9f28-c16f07ae010c.pdf?c=1632337061>

[As per journal policy, preprint can be published as a journal article, provided it is not published in any other journal]

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ETHICAL APPROVAL

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee".

The care and handling of animals complied with the National Institutes of Health Manual on the care and use of laboratory animals (National Research Council, 2010) and was approved by the Animal Use Ethics Committee of the Landmark University, Omu-Aran, Nigeria, with the approval number LMU/EC/098/2019. Furthermore, this study was conducted according to the ARRIVE guidelines (Percie du Sert et al., 2020).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abdelkader, N. F., Eitah, H. E., Maklad, Y. A., Gamaleldin, A. A., Badawi, M. A., & Kenawy, S. A. (2020). New combination therapy of gliclazide and quercetin for protection against STZ-induced diabetic rats. *Life Sciences*, *247*, 117458.
- Ajiboye, B. O., Ojo, O. A., Akuboh, O. S., Abiola, O. M., Idowu, O., & Amuzat, A. O. (2018). Anti-hyperglycemic and anti-inflammatory activities of polyphenolic-rich extract of *Syzygium cumini* Linn leaves in alloxan-induced diabetic rats. *Journal of Evidence-Based Integrative Medicine*, *23*, 2515690X18770630.
- Akdis, C. A., Joss, A., Akdis, M., Faith, A., & Blaser, K. (2000). A molecular basis for T cell suppression by IL-10: CD28-associated IL-10 receptor inhibits CD28 tyrosine phosphorylation and phosphatidylinositol 3-kinase binding. *The FASEB Journal*, *14*(12), 1666-1668.
- Berryman, A. M., Maritim, A. C., Sanders, R. A., & Watkins, J. B. (2005). Influence of treatment of diabetic rats with combinations of pycnogenol, β -carotene, and α -lipoic acid on parameters of oxidative stress. *Journal of Biochemical and Molecular Toxicology*, *18*(6), 345-352.
- Domingueti, C. P., Dusse, L. M., das Graças Carvalho, M., de Sousa, L. P., Gomes, K. B., & Fernandes, A. P. (2016). Diabetes mellitus: The linkage between oxidative stress, inflammation, hypercoagulability, and vascular complications. *Journal of Diabetes and its Complications*, *30*(4), 738-745.
- Domingueti, C. P., Fóscolo, R. B., Dusse, L. M., Reis, J. S., Carvalho, M. D., Gomes, K. B., & Fernandes, A. P. (2018). Association of different biomarkers of renal function with D-dimer levels in patients with type 1 diabetes mellitus. *Archives of Endocrinology and Metabolism*, *62*, 27-33.
- Gratas-Delamarche, A., Derbré, F., Vincent, S., & Cillard, J. (2014). Physical inactivity, insulin resistance, and the oxidative-inflammatory loop. *Free Radical Research*, *48*(1), 93-108.
- Macdonald, I., Mohammed, A., & Adeboye, A. (2017). Alloxan-induced diabetes, a common model for evaluating the glycemic-control potential of therapeutic compounds and plant extracts in experimental studies. *Medicina*, *53*(6), 365-374.
- Nayak, S. S., & Pattabiraman, T. N. (1981). A new colorimetric method for the estimation of glycosylated hemoglobin. *Clinica Chimica Acta*, *109*(3), 267-274.
- Perez Gutierrez, R. M., Flores Cotera, L. B., & Gonzalez, A. M. (2012). Evaluation of the antioxidant and anti-glycation effects of the hexane extract from *Piper auritum* leaves in vitro and beneficial activity on oxidative stress and advanced glycation end-product-mediated renal injury in streptozotocin-treated diabetic rats. *Molecules*, *17*(10), 11897-11919.
- Rodrigues, B., Poucheret, P., Battell, M. L., & McNeill, J. H. (2018). Streptozotocin-induced diabetes: Induction, mechanisms, and dose dependency. In *Experimental models of diabetes* (pp. 3-17). Routledge.

- Rosero, J. C., Cruz, S., Osorio, C., & Hurtado, N. (2019). Analysis of phenolic composition of byproducts (seeds and peels) of avocado (*Persea americana* Mill.) cultivated in Colombia. *Molecules*, 24(17), 3209.
- Rouhi, Y., Naseri, F., Tahmasebi, E., Tebyanian, H., & Moghaddam, M. M. (2020). Effect of vitamin D on the expression level of the IL-10 and IL-4 in asthmatic mice. *Biointerface Research in Applied Chemistry*, 10(6), 7077-7084.
- Shahwan, M., Alhumaydhi, F., Ashraf, G. M., Hasan, P. M., & Shamsi, A. (2022). Role of polyphenols in combating type 2 diabetes and insulin resistance. *International Journal of Biological Macromolecules*.
- Sindhughosa, D. A., & Pranamartha, A. G. (2017). The involvement of proinflammatory cytokines in diabetic nephropathy: Focus on interleukin 1 (IL-1), interleukin 6 (IL-6), and tumor necrosis factor-alpha (TNF- α) signaling mechanism. *Bali Medical Journal*, 6(1), 44-51.
- Wada, J., & Makino, H. (2013). Inflammation and the pathogenesis of diabetic nephropathy. *Clinical Science*, 124(3), 139-152.
- Yang, D. K., & Kang, H. S. (2018). Anti-diabetic effect of cotreatment with quercetin and resveratrol in streptozotocin-induced diabetic rats. *Biomolecules & Therapeutics*, 26(2), 130.
- Yang, J., Ma, Y., Li, T., Pang, Y., Zhang, H., Xie, Y., Liu, H., Sun, Y., Ren, J., & Jin, J. (2023). Ameliorative effects of *Bifidobacterium animalis* subsp. *lactis* J-12 on hyperglycemia in pregnancy and pregnancy outcomes in a high-fat-diet/streptozotocin-induced rat model. *Nutrients*, 15(1), 170.
- Yaribeygi, H., Butler, A. E., Barreto, G. E., & Sahebkar, A. (2019). Antioxidative potential of antidiabetic agents: A possible protective mechanism against vascular complications in diabetic patients. *Journal of Cellular Physiology*, 234(3), 2436-2446.
- Zafar, H., Sheikh, M. A., Hussain, F., & Maan, M. A. (2012). Inhibition of protein glycation and advanced glycation end products by ascorbic acid. *African Journal of Biotechnology*, 11(51), 11309-11314.
- Alkhalaf, M. I., Alansari, W. S., Ibrahim, E. A., & Elhalwagy, M. E. (2019). Anti-oxidant, anti-inflammatory and anti-cancer activities of avocado (*Persea americana*) fruit and seed extract. *Journal of King Saud University - Science*, 31(4), 1358-1362.
- Bigagli, E., & Lodovici, M. (2019). Circulating oxidative stress biomarkers in clinical studies on type 2 diabetes and its complications. *Oxidative Medicine and Cellular Longevity*, 2019.
- Colombo, R., & Papetti, A. (2019). Avocado (*Persea americana* Mill.) by-products and their impact: From bioactive compounds to biomass energy and sorbent material for removing contaminants. *International Journal of Food Science & Technology*, 54(4), 943-951.
- Figuroa, J. G., Borrás-Linares, I., Lozano-Sánchez, J., & Segura-Carretero, A. (2018). Comprehensive characterization of phenolic and other polar compounds in the seed and seed coat of avocado by HPLC-DAD-ESI-QTOF-MS. *Food Research International*, 105, 752-763.
- National Research Council. (2010). *Guide for the care and use of laboratory animals*. National Academies Press.
- Oguntibeju, O. O. (2019). Hypoglycaemic and anti-diabetic activity of selected African medicinal plants. *International Journal of Physiology Pathophysiology Pharmacology*, 11(6), 224.
- Oluba, O. M., Adebisi, F. D., Dada, A. A., Ajayi, A. A., Adebisi, K. E., Josiah, S. J., & Odotuga, A. A. (2019). Effects of *Talinum triangulare* leaf flavonoid extract on streptozotocin-induced hyperglycemia and associated complications in rats. *Food Science & Nutrition*, 7(2), 385-394.
- Oluba, O. M., Akpor, O. B., Adebisi, F. D., Josiah, S. J., Alabi, O. O., Shoyombo, A. O., & Olusola, A. O. (2020). Effects of co-administration of *Ganoderma* terpenoid extract with chloroquine on inflammatory markers and antioxidant status in *Plasmodium berghei*-infected mice. *Journal of Integrative Medicine*, 18(6), 522-529.
- Özcan, M. M., Al Juhaimi, F., Ahmed, I. A., Uslu, N., Babiker, E. E., & Ghafoor, K. (2020). Effect of microwave and oven drying processes on antioxidant activity, total phenol and phenolic compounds of kiwi and pepino fruits. *Journal of Food Science and Technology*, 57, 233-242.
- Pamplona, R. (2008). Membrane phospholipids, lipoxidative damage and molecular integrity: A causal role in aging and longevity. *Biochimica et Biophysica Acta - Bioenergetics*, 1777(10), 1249-1262.
- Percie du Sert, N., Hurst, V., Ahluwalia, A., Alam, S., Avey, M. T., Baker, M., Browne, W. J.,

- Clark, A., Cuthill, I. C., Dirnagl, U., & Emerson, M. (2020). The ARRIVE guidelines 2.0: Updated guidelines for reporting animal research. *Journal of Cerebral Blood Flow & Metabolism*, 40(9), 1769-1777.
- Simić, V. M., Rajković, K. M., Stojičević, S. S., Veličković, D. T., Nikolić, N. Č., Lazić, M. L., & Karabegović, I. T. (2016). Optimization of microwave-assisted extraction of total polyphenolic compounds from chokeberries by response surface methodology and artificial neural network. *Separation and Purification Technology*, 160, 89-97.
- Vinha, A. F., Sousa, C., Soares, M. O., & Barreira, S. V. (2020). Avocado and its by-products: Natural sources of nutrients, phytochemical compounds and functional properties. *Current Research in Agricultural and Food Science*, 1, 82-96.
- Wu, C. H., Huang, S. M., Lin, J. A., & Yen, G. C. (2011). Inhibition of advanced glycation end-product formation by foodstuffs. *Food Function*, 2(5), 224-234.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/127435>