



Laparoscopic Repair of Ileal Perforation Post Blunt Abdominal Trauma: A Case Report

**Mohamed M Elgeldawy ^{a*}, Reham F Alfalakawi ^a,
Fahad Alabhoul ^a and Ali H Dokhi ^a**

^a Department of Surgery, Al Adan Hospital, Ministry of Health, Kuwait.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajcrs/2025/v8i1591>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129224>

Case Report

Received: 02/11/2024

Accepted: 05/01/2025

Published: 11/01/2025

ABSTRACT

Blunt trauma to the abdomen is considered one of the most leading causes of death among all age groups. Although the outcome of patients with blunt abdominal injury has improved in the last two decades, in patients with multiple organ injuries, the in-hospital mortality rate was reported as 3–10%. Delayed diagnosis of small bowel perforation can result in serious complications and death, making timely diagnosis critical in the treatment of these patients. The diagnosis of small bowel perforation is usually made by imaging techniques that detect the presence of fluids or gases in the peritoneum or mediastinum. There are different treatment options for small bowel perforation; conservative treatment is not favoured.

Herein, we present the case of a 35-year-old gentleman diagnosed with a perforated viscus following a road traffic accident. A diagnostic laparoscopy was performed, and a 1 cm perforation of

*Corresponding author: E-mail: thesurgeon76@yahoo.com;

the terminal ileum was repaired. Postoperative recovery was uneventful except for mild paralytic ileus. The patient was discharged 8 days later and reviewed at the surgical outpatient department after one- and two-months post-surgery, with no reported complaints.

Keywords: Blunt abdominal trauma; Ileal perforation; laparoscopic repair; Road traffic accident.

1. INTRODUCTION

“Blunt trauma to the abdomen is considered one of the most leading causes of death among all age groups. Thus, it is an important topic of discussion and research” (Dauterive et al., 1985). “Although the outcome of patients with blunt abdominal injury has improved in the last two decades, in patients with multiple organ injuries, the in-hospital mortality rate was reported as 3–10%” (Mirzamohamadi et al., 2024). There are many causes of blunt abdominal trauma (BAT) such as assault, falls from height, and animal attacks. However, road traffic accidents (RTA) are considered the most common cause of BAT to the abdomen (Hughes and Elton, 2002, Kurane and Ugane, 2017). “The diagnosis of small bowel perforation in patients with BAT is infrequent as the injury needs to be severe to cause perforation. A hollow viscus injury is not usually suspected unless the clinical condition is highly suggestive” (Kurane and Ugane, 2017). “A delay in the diagnosis and treatment of the hollow viscus injury will cause the development of peritonitis, and hemodynamic instability leading to increased mortality and morbidity” (Hughes and Elton, 2002). “Although small bowel perforation is rare, it is a clinically complex medical condition with high mortality and should be diagnosed and managed early for a better prognosis” (Onken et al., 2025). Properly managing small bowel perforation can go through early stabilisation, administration of antibiotics, and surgical source control, with prompt operative intervention (Fakhry et al., 2019).

2. CASE REPORT

A 35-year-old gentleman presented to the emergency room of the casualty department in an emergency center away from our hospital by about 80 Kilometers at 2:00 am following RTA, he was complaining of right-side abdominal pain. He was vitally stable, and his abdominal examination showed right-sided abdominal tenderness. His labs were normal (WBCs 8.5, Hb 12.4). He was managed according to the ATLS protocol. His chest and pelvic X-rays were normal. Pan computed tomography (PAN-CT) (brain, chest abdomen and pelvic) was requested, which revealed minimal pelvic free fluid with mild ascending colon oedema (Fig. 1). At 5:00 am, the patient was admitted to our hospital for observation. A serial abdominal examination was started, but the abdominal examination revealed board-like rigidity at 9:00 am. The patient was shifted to the operating theatre at 9:30 am as an erect chest x-ray revealed massive air under the diaphragm (Fig. 2). Diagnostic laparoscopy was started, and we found intestinal contents inside the peritoneal cavity then running of the small bowel was done and revealed a 1 cm perforation at the ileum, about 120 cm from the ileocaecal junction (Fig. 3). Laparoscopic repair was done by Vicryl 2/0 in 2 layers and generous peritoneal washing with warm saline and 2 drains were fixed one in the pelvis and one subdiaphragmatic above the liver. post-operative recovery was complicated by paralytic ileus, and the patient was discharged after 8 days. After 1 month and 2 months, the patient was reviewed in the surgical outpatient clinic, and he has no complaints.




Fig. 1. CT abdomen, NO pneumoperitoneum was detected




Fig. 2. ERECT CHEST X-RAY showing bilaterally massive pneumoperitoneum




Fig. 3. Intraoperative finding of intestinal contents and the perforation of the ileum

3. DISCUSSION

Because of modern life, developments in industry, various transport systems and violence, there is an increase in the number of patients with BAT (Bouzat et al., 2020). The bowel injury following BAT is frequently undetected, so the diagnosis is usually too late despite advanced imaging techniques such as focused abdominal sonography for trauma and computed tomography scans (Yavaş et al., 2023). Apart from our patient, he was diagnosed within hours and was in the operating theatre within about 7-8 hours. Traumatic causes of small bowel perforation can be due to blunt or penetrating injuries (our case was after BAT) (Aljehaiman et al., 2024). The small bowel occupies the majority of the abdominal cavity, making it the most frequently injured hollow organ in the abdomen, particularly in cases of penetrating trauma such as stab wounds or gunshots. Blunt abdominal trauma due to RTA is one of the most common causes of small bowel perforation (as seen in our case). Seat-belt users account for approximately 75% of gastrointestinal trauma with potential

perforation (Aljehaiman et al., 2024). There are two types of small bowel perforation: free or contained perforation. Free perforations create an open hole that allows the intestinal material to escape into the normally sterile peritoneal cavity and can cause infection, inflammation, and other complications to the surrounding tissues and organs (as you can see in our case). On the other hand, contained perforations are those in which an ulcer creates a full-thickness hole, but the adjacent organs seal it and prevent the free spillage of the intestinal content (Hafner et al., 2023). "A high degree of suspicion of intra-abdominal injuries, even in cases following minor trauma will prevent the diagnostic errors. Early diagnosis and effective management of gastrointestinal perforations (as revealed in our case) following BAT to the abdomen helps in decreasing overall morbidity and mortality among these patients" (Gurav et al., 2024). "The symptoms of small bowel perforation may range from localised abdominal pain to systemic symptoms like fever, nausea, vomiting, and even shock. Additionally, rigidity, guarding, and other peritoneal signs which may appear later"

(Katwal et al., 2023). “Although erect plain X-rays are usually the first imaging option to start with and can detect signs of perforation, they cannot localise the site of perforation” (Kulinna-Cosentini et al., 2024). The abdominal computed tomography scan is regarded as the imaging modality of choice for hemodynamically stable patients; however, negative results don’t rule out the diagnosis of small bowel perforation (Bini et al., 2020) (as in our case, **NO** pneumoperitoneum in the abdominal computed tomography was detected). In the **EAST trial**, 13% of the study samples who were discovered to have a small bowel perforation at laparotomy had no positive findings before surgery (Loftus et al., 2019). “Delays in the diagnosis of small bowel perforation can result in serious complications and death, making timely diagnosis critical in the treatment of these patients” (Fakhry et al., 2019). In trauma patients, laparoscopic interventions with normal haemodynamic parameters are an excellent modality to identify any abdominal injuries. Laparoscopy is more efficient and cost-effective and associated with fewer complications in comparison with traditional laparotomies (El-Bendary et al., 2016). Although there are different options of treatments for the perforation of the small bowel, conservative treatment is not favoured (Pouli et al., 2020). Laparoscopy and open surgery (primary repair, anastomosis, and stoma) are mainly used, which may be related to timely preoperative management, good resuscitation of septic shock, and timely application of broad-spectrum antibiotics in the perioperative period (Li et al., 2023). The mortality rates among patients diagnosed with BAT with small bowel perforation are linked to the delay in initial diagnosis, development of multi-organ failure, and sepsis as emphasized by the **EAST trial**, which reported a rise in mortality rate from 4% to 16% when the diagnosis is delayed for more than 24h. it was further highlighted that when small bowel perforation was present the mortality rate rose from 14% to 19% (Staab et al., 2024).

4. CONCLUSION

Small bowel perforation following blunt abdominal trauma is rare. Thorough clinical assessment through serial abdominal examinations, alongside investigations that may sometimes be inconclusive, and timely surgical intervention can play a crucial role in reducing morbidity and mortality. The minimally invasive techniques such as laparoscopy offer an

excellent option for the effective management of blunt abdominal trauma.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

CONSENT

Written informed consent was obtained from the patient for publication of this case report and accompanying images.

ETHICAL APPROVAL

The author(s) have collected and preserved written ethical approval as per international or university standards.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Aljehaiman, F., Almalki, F. J., Alhusain, A., Alsalamah, F., Alzahrani, K., Alharbi, A., & Alkhulaiwi, H. (2024). Prevalence, pattern, mortality, and morbidity of traumatic small bowel perforation at King Abdulaziz Medical City: A retrospective cohort study. *Cureus*, 16(1), e52313.
- Bini, R., Ronchetta, C., Picotto, S., Scozzari, G., Gupta, S., Frassini, S., & Chiara, O. (2020). Importance of CT-scan predicting clinical outcomes in gastrointestinal perforation. *Annals of Translational Medicine*, 8(21).
- Bouzat, P., Valdenaire, G., Gauss, T., Charbit, J., Arvieux, C., Balandraud, P., Bobbia, X., David, J. S., Frandon, J., Garrigue, D., & Long, J. A. (2020). Early management of severe abdominal trauma. *Anaesthesia Critical Care & Pain Medicine*, 39(2), 269–277.
- Dauterive, A. H., Flancbaum, L., & Cox, E. F. (1985). Blunt intestinal trauma: A modern-day review. *Annals of Surgery*, 201(2), 198–203.
- El-Bendary, Y. B., Al-Adawi, J., & Al-Qadhi, H. (2016). The use of laparoscopy in the

- management of trauma patients: Brief review. *Sultan Qaboos University Medical Journal*, 16(1), e9–e14.
- Fakhry, S. M., Allawi, A., Ferguson, P. L., Michetti, C. P., Newcomb, A. B., Liu, C., & Brownstein, M. R. (2019). Blunt small bowel perforation (SBP): An Eastern Association for the Surgery of Trauma multicenter update 15 years later. *Journal of Trauma and Acute Care Surgery*, 86, 642–650.
- Fakhry, S. M., Allawi, A., Ferguson, P. L., Michetti, C. P., Newcomb, A. B., Liu, C., & Brownstein, M. R. (2019). Blunt small bowel perforation (SBP): An Eastern Association for the Surgery of Trauma multicenter update 15 years later. *Journal of Trauma and Acute Care Surgery*, 86(4), 642–650.
- Gurav, S. P., Quraishi, A. H., Umare, G., & Tongse, P. S. (2024). A prospective observational study of gastrointestinal perforations following blunt trauma abdomen. *International Surgery Journal*, 11(5), 737–743.
- Hafner, J., Tuma, F., Hoilat, G. J., & Marar, O. (2023). Intestinal perforation. In *StatPearls* [Internet]. StatPearls Publishing.
- Hughes, T. M. D., & Elton, C. (2002). The pathophysiology and management of bowel and mesenteric injuries due to blunt trauma. *Annals of Surgery*, 33, 120–128.
- Katwal, S., Ghimire, A., & Joshi, N. (2023). Uncommon presentation: Isolated small bowel perforation after blunt abdominal trauma in a pediatric patient: A case report. *Annals of Medicine and Surgery*, 85(12), 6222–6226.
- Kulinna-Cosentini, C., Hodge, J. C., & Basalamah, A. (2024). The role of radiology in diagnosing gastrointestinal tract perforation. *Best Practice & Research Clinical Gastroenterology*, 101928.
- Kurane, S. B., & Ugane, S. P. (2017). A clinical study of hollow viscus injury due to blunt trauma abdomen. *International Journal of Research in Medical Sciences*, 5(11), 5017.
- Li, Z., Li, S., Wu, C., Liu, Z., Zhao, Y., Luan, J., Miao, S., & Wang, P. (2023). Diagnostic and therapeutic procedures of small bowel perforation: A retrospective monocentric cohort study. *World Journal of Surgical Infection*, 2(1), 7–11.
- Loftus, T. J., Morrow, M. L., Lottenberg, L., Rosenthal, M. D., Croft, C. A., Smith, R. S., Moore, F. A., Brakenridge, S. C., Borrego, R., Efron, P. A., & Mohr, A. M. (2019). Occult bowel injury after blunt abdominal trauma. *American Journal of Surgery*, 218(2), 266–270.
- Mirzamohamadi, S., HajiAbbasi, M. N., Baigi, V., Salamati, P., et al. (2024). Patterns and outcomes of patients with abdominal injury: A multicenter study from Iran. *BMC Emergency Medicine*, 24(1), 91.
- Onken, F., Senne, M., Königsrainer, A., & Wichmann, D. (2022). Classification and treatment algorithm of small bowel perforations based on a ten-year retrospective analysis. *Journal of Clinical Medicine*, 11, 5748. <https://doi.org/10.3390/jcm11195748>
- Pouli, S., Kozana, A., Papakitsou, I., Daskalogiannaki, M., & Raissaki, M. (2020). Gastrointestinal perforation: Clinical and MDCT clues for identification of aetiology. *Insights into Imaging*, 11.
- Staab, V., Naganathan, S., McGuire, M., Pinto, J. M., & Pall, H. (2024). Gastrointestinal perforation with blunt abdominal trauma in children. *Children (Basel)*, 11(6), 612.
- Yavaş, Ö., Çelik, A., Altuntaş, G., Altuntaş, M., Bilir, Ö., & Ersunan, G. (2023). The role of repeated extended FAST in patients with stable blunt thoracoabdominal trauma. *Turkish Journal of Trauma & Emergency Surgery/Ulusal Travma ve Acil Cerrahi Dergisi*, 29(5).

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/129224>